



MP Biomedicals, LLC.

29525 Fountain Parkway  
Solon, Ohio 44139

Telephone: 440/337-1200  
Toll Free: 800/854-0530  
Fax: 440/337-1180  
mailto:biotech@mpbio.com  
web: www.mpbio.com

# cappel

**Anti-Desmin II Monoclonal 1.0 ML**

**Catalog #: 10519**

**Lot #: 8575C**

**Species:** Mouse Monoclonal; Hybridoma: mouse myeloma cells SP2/0 Ag14-BALB/c.

**Clone:** D9

**Ig** **Class:** IgG<sub>1</sub>

**Reagent:** Culture supernatant

**Immunogen:** Human leiomyoma cell isolate

**Specificity:** Antibody reacts with the 53 kD desmin protein. Immunoblotting: protein.

**Cross-Reactivity:** Most With: mammals.

**Localization:** Cytoplasm

**Application:** Frozen Sections/Cell Preparations; paraffin sections (no pretreatment).

**Note:** Frozen Sections; fixation with acetone for 5 minutes at 20°C. Paraffin Sections; fixation with formaldehyde.

**Positive Control:** Skeletal and smooth muscles myosarcomas

**Dilution:** 1:40 (Indirect method, 30 minute incubation)  
Dilute immediately before use with PBS. Optimal dilution should be tested by serial dilution.

**Storage:** Store undiluted antibody at 2-8°C. For prolonged storage, store aliquots at -20°C.

**Reference:** 1. Muyen G.N.P. van et al. Coexpression of Intermediate Filament Polypeptides in Fetal and Adult Tissues (1987). **Lab. Invest.**, 57, 4, 359-369.

This product is intended for Research Use Only, "for Tests *in vitro*", not for use in diagnostic or therapeutic procedures involving humans or animals.

This product contains sodium azide. To prevent formation of toxic vapors, do not mix with strongly acidic solutions. To prevent formation of potentially explosive metallic azides in metal plumbing, always wash into drain with copious quantities of water.  
R0703

  
Approved by: Joseph Dietz, Ph.D.  
Quality Control Director

**TO PLACE AN ORDER IN THE U.S., PLEASE CALL 800-854-0530**  
**OR TO OBTAIN TECHNICAL ASSISTANCE IN THE U.S., PLEASE CALL 800-279-5490**  
**OUTSIDE THE U.S., CONTACT YOUR LOCAL DISTRIBUTOR.**