



MP Biomedicals, LLC

29525 Fountain Parkway
Solon, Ohio 44139

Telephone: 440/337-1200

Toll Free: 800/854-0530

Fax: 440/337-1180

mailto:biotech@mpbio.com

web: www.mpbio.com

cappel

**GOAT AFFINITY PURIFIED ANTIBODY TO
HUMAN IGA (alpha CHAIN) - 1MG**

Catalog #: 55068

Lot #: 07993

Total Protein: 1.02 mg/ml
Active Antibody: 1.0 mg/ml

DESCRIPTION:

Product is goat affinity purified antibody to human IgA (α chain) and buffer salts.

The affinity purified antibody is prepared from the specific goat IgG fraction by solid phase absorption. The resulting affinity purified antibody is dialyzed into 0.02M sodium phosphate, 0.14M sodium chloride, pH 7.3, adjusted to standard titer, filtered through a 0.22 μ m filter, and vialled. No preservative has been added.

USES:

This product is suitable for use as a precipitating antibody. It gives sharp precipitin lines in immunoelectrophoresis and immunodiffusion techniques.

This product is suitable for conjugation with enzymes, radiolabels, fluorochromes or other markers, and for attachment to solid supports for use as an immunoabsorbant.

This product is suitable for use as a primary reagent in enzyme immunoassays, fluorescein immunoassays and blot, cell or tissue immunostaining. These affinity purified antibodies are recommended to avoid non-specific binding from the inherent antibodies of host animals. (Note: F(ab')₂ fragments are recommended for staining of cells or tissues which contain Fc receptors. See **Related Product Ordering Information** for these products.)

SUGGESTED WORKING DILUTIONS:

- 1: 50 - 1: 200 for Tissue Staining
- 1: 100 - 1: 500 for Immunoblotting
- 1: 400 - 1: 2,000 for Elisa

It is recommended that each lab obtain their own optimal working dilution for each assay

SPECIFICATIONS:

The total protein is measured by absorbance (A_{280nm}). Antibody titer is standardized by equivalence-point precipitation. Each affinity purified antibody is tested for purity at 10 mg/ml using immunoelectrophoresis. The product is goat IgG; no trace of albumin is detected. It shows reactivity to human IgA; cross-reactivity to other species may exist. Antibody activity to IgG, IgM, or non-immunoglobulin serum proteins is not present.

EXPIRATION DATE:

2/28/2021

STORAGE:

Store unopened product at 2-8°C. For long term storage, aliquot and store with 40% glycerol (v/v) at -20°C.

WARNING AND CAUTION:

This product is intended **FOR RESEARCH USE ONLY**, not for use in diagnostic or therapeutic procedures involving humans or animals.

RELATED PRODUCT ORDERING INFORMATION:

| <u>Catalog #</u> | <u>Description</u> | <u>Size</u> |
|------------------|--|-------------|
| 55082 | GOAT ANTI-HUMAN IGA (□ CHAIN) | 2ML |
| 55003 | IGG GOAT ANTI-HUMAN IGA (□ CHAIN) | 5ML |
| 55047 | F(AB') ₂ GOAT ANTI-HUMAN IGA (□ CHAIN) | 2ML |
| 55077 | FITC GOAT ANTI-HUMAN IGA (□ CHAIN) | 2ML |
| 55178 | FITC F(AB') ₂ GOAT ANTI-HUMAN IGA (□ CHAIN) | 2ML |
| 55195 | FITC AFF.PUR. GOAT ANTI-HUMAN IGA (□ CHAIN) | 2MG |
| 55216 | HRP GOAT ANTI-HUMAN IGA (□ CHAIN) | 2ML |
| 55243 | HRP F(AB') ₂ GOAT ANTI-HUMAN IGA (□ CHAIN) | 2ML |
| 59286 | AKP AFF.PUR. GOAT ANTI-HUMAN IGA (□ CHAIN) | 1ML |
| 55906 | PURIFIED HUMAN SERUM IGA | 5MG |



Approved by: *John Huang, PhD*
Quality Control Manager

TO PLACE AN ORDER IN THE U.S., PLEASE CALL 800-854-0530
OR TO OBTAIN TECHNICAL ASSISTANCE IN THE U.S., PLEASE CALL 800-279-5490
OUTSIDE THE U.S., CONTACT YOUR LOCAL DISTRIBUTOR.