



MP Biomedicals, LLC.

29525 Fountain Parkway
Solon, Ohio 44139

Telephone: 440/337-1200
Toll Free: 800/854-0530
Fax: 440/337-1180
mailto:biotech@mpbio.com
web: www.mpbio.com

cappel

**GOAT ANTISERUM TO
HUMAN FIBRONECTIN - 2ML**

Catalog #: 55125

Lot #: 00327

Total Protein: 42.1 mg/ml

DESCRIPTION:

Product is the lyophilized powder of goat antiserum to human fibronectin and buffer salts.

The antiserum is prepared by hyperimmunizing goats with highly purified human fibronectin; antiserum is delipidated and made specific by solid-phase absorption. The resulting antiserum is dialyzed into 0.02M sodium phosphate, 0.14M sodium chloride, pH 7.3, with 0.05% sodium azide, adjusted to standard titer, filtered through a 0.22 μ m filter, vialled and lyophilized.

USES:

This product is suitable for use as a precipitating antibody. It gives sharp precipitin lines in immunoelectrophoresis and immunodiffusion techniques.

(Note: IgG fractions are suitable for conjugation with enzymes, radiolabels, fluorochromes or other markers and for attachment to solid supports for use as an immunoabsorbant. F(ab')₂ fragments are recommended for staining of cells or tissues which contain Fc receptors. See **Related Product Ordering Information** for these products.)

SUGGESTED WORKING DILUTIONS:

1:50 - 1:200 for Tissue Staining
1:100 - 1:500 for Immunoblotting
1:400 - 1:2,000 for ELISA

It is recommended that each lab obtain their own optimal working dilution for each assay.

SPECIFICATIONS:

The total protein is measured using the Biuret procedure, with bovine albumin as a standard. Antibody titer is standardized with an in-house control by immunoelectrophoresis. Each antiserum is tested for specificity at a minimum of 80 mg/ml using immunoelectrophoresis. It shows reactivity to human fibronectin; cross-reactivity to other species may exist. Antibody activity to human fibrinogen or other serum proteins is not present.

RECONSTITUTION:

Reconstitute product with 2.0 ml of deionized or distilled water. Gentle swirling may be used to speed rehydration. Avoid vigorous shaking of the reconstituted material.

STORAGE:

Store lyophilized product at 2-8°C. Store reconstituted material no longer than 4 weeks at 2-8°C. For long-term storage, aliquot and freeze reconstituted product at -20°C or lower.

WARNING AND CAUTION:

This product is intended **FOR RESEARCH USE ONLY**, not for use in diagnostic or therapeutic procedures involving humans or animals.

This product contains sodium azide. To prevent formation of toxic vapors, do not mix with strongly acidic solutions. To prevent formation of potentially explosive metallic azides in metal plumbing, always wash into drain with copious quantities of water.

RELATED PRODUCT ORDERING INFORMATION:

Catalog # Description Size

55037 (02010831)	IGG GOAT ANTI-HUMAN FIBRONECTIN	5ML
55065 (03010831)	F(AB') ₂ GOAT ANTI-HUMAN FIBRONECTIN	2ML
55913 (60010830)	PURIFIED HUMAN FIBRONECTIN	1MG
55170 (12010831)	FITC GOAT ANTI-HUMAN FIBRONECTIN	2ML
55193 (13010831)	FITC F(AB') ₂ GOAT ANTI-HUMAN FIBRONECTIN	2ML
55240 (32010831)	HRP GOAT ANTI-HUMAN FIBRONECTIN	2ML


Approved by Joseph Dietz, Ph.D.
Quality Control Director

TO PLACE AN ORDER IN THE U.S., PLEASE CALL 800-854-0530
OR TO OBTAIN TECHNICAL ASSISTANCE IN THE U.S., PLEASE CALL 800-279-5490
OUTSIDE THE U.S., CONTACT YOUR LOCAL DISTRIBUTOR.